Bariatric Phase 2 Full Liquid Diet

Breakfast

Instructions:

Drink 1-2 oz. liquid every 10-15 minutes. Do not drink 30 minutes before eating.

Do not drink 30 minutes before eating.				
<u>Time</u>	Menu	Cal	Pro	Carbs
7:00am	½ - ¾ cup decaf coffee/tea	0	0	0
8:00am	SF popsicle	0	0	0
9:00 am	½ cup HP skim milk (1 sc)	65	10	6
10:00am	½ cup diluted apple juice	30	0	7
	~16 fluid oz.	105 Cal	10gP	13g

Sugar Sub.

4oz high protein supplement

16 oz. bottle of Water will be provided daily.

NAME: ______ ROOM #____ Daily menus provides: 75 cal 0g Pro 18g Carbs

Bariatric Phase 2 Full Liquid Diet

Lunch

Instructions:

Drink 1-2 oz. liquid every 10-15 minutes. Do not drink 30 minutes before eating.

DO HOU	Do not drink 30 initiales before eating.			
<u>Time</u>	<u>Menu</u>	<u>Cal</u>	<u>Pro</u>	<u>Carbs</u>
12:00pm	½ - ¾ cup decaf coffee/tea	0	0	0
2:00pm	unsweetened HP applesauce (2 sc)	100	12	13.5
3:00pm	SF popsicle	0	0	0
4:00pm	½ - ¾ cup water	0	0	0
	~18 fluid oz.	100 Cal	12gF	13.5g

Sugar Sub.

4oz high protein supplement

NAME: ROOM

Bariatric Phase 2 Full Liquid Diet

Dinner

Instructions:

Drink 1-2 oz. liquid every 10-15 minutes. Do not drink 30 minutes before eating.

Do not drink 30 minutes before eating.				
<u>Time</u>	<u>Menu</u>	<u>Cal</u>	<u>Pro</u>	Carb
6:00pm	½ - ¾ cup decaf coffee/tea	0	0	0
7:00pm	½ cup HP soup (1 sc)	25	6	0
8:00pm	6 oz Dannon Light Fit 'n Creamy Yogurt®	80	5	15
9:00pm	½ - ¾ cup water	0	0	0
	~ 20 fluid oz. 105	5Cal 1	1gP	15g

Sugar Sub.

4 oz. high protein supplement

NAME:	ROOM #